

Working Guidelines for Breast Imaging in Relation to COVID-19 Vaccination

- **Unilateral axillary lymphadenopathy on side of COVID Vaccination**
 - **Approximately 15% of patients**
 - **starting 2-4 days after vaccination and lasting 4-6 weeks.**
 - **Lymph node enlargement can be seen on mammography and MRI, leading to additional ultrasound follow-up, and possible biopsy.**

Working Guidelines for Breast Imaging in Relation to COVID-19 Vaccination

SCHEDULING SCREENING MAMMOGRAMS AND MRI:

1. If the patient has already had the COVID-19 vaccine,
 - schedule screening mammograms or MRI 4-6 weeks after the 2nd vaccine dose to avoid false positive results.
2. If the patient is to be scheduled for the vaccine
 - try to have the screening mammogram or MRI performed before the vaccine is given.

FOR PATIENTS WITH A DIAGNOSTIC BREAST FINDINGS

1. Patients with breast complaints and axillary lump/tenderness/pain should undergo diagnostic evaluation regardless of vaccination status. If an enlarged axillary lymph node is found, ultrasound will be performed.
 - If the enlarged node is on the side of COVID vaccination within the past 4 weeks, and if clinically appropriate, follow-up ultrasound will be performed in 4-6 weeks since the overwhelming majority of the enlarged lymph nodes will resolve within that time frame.
 - With persistent enlargement or increase size, biopsy may be necessary.


Working Guidelines for Breast Imaging in Relation to COVID-19 Vaccination

Attached is a link for more comprehensive discussion provided by the Society of Breast Imaging.

<https://www.sbi-online.org/Portals/0/Position%20Statements/2021/SBI-recommendations-for-managing-axillary-adenopathy-post-COVID-vaccination.pdf>

We continue to recommend annual screening mammograms for all women of average risk after age 40 based on the recommendations of The American College of Radiology and The National Comprehensive Care Network (NCCN).

Working Guidelines for Breast Imaging in Relation to COVID-19 Vaccination



Working Guidelines for Breast Imaging in Relation to COVID-19 Vaccination

SBI Screening Mammography Recommendations for Women Receiving the COVID-19 Vaccine



Screening mammograms and COVID-19 vaccines are both very important for your health.

■ COVID-19 vaccine

Some women who receive the COVID-19 vaccine develop swollen lymph nodes under their arm on the same side as their vaccine injection. This is the normal immune reaction to a vaccine. These swollen lymph nodes usually return to normal on their own in a few days or weeks.

■ Why do swollen lymph nodes matter?

Breast radiologists look closely for any changes on your mammogram. Swollen lymph nodes under one arm can be seen on a mammogram and can be a rare sign of breast cancer.

■ What happens if there are swollen lymph nodes on my mammogram?

Depending on your medical history and when you received your vaccine, the breast radiologist may recommend that you return to the breast center for an ultrasound of your underarm area and they also may recommend a follow up exam to show that the lymph nodes have returned to normal size.

■ When should I schedule my screening mammogram?

Try to schedule your screening mammogram before your first COVID-19 vaccine dose or at least 4 weeks after your second vaccine dose. This reduces the chance that swollen lymph nodes from the vaccine will appear on your mammogram.

■ What if my mammogram is already scheduled?

Keep your vaccination appointment. Getting vaccinated is critical to stop the spread of COVID-19. Consider rescheduling your screening mammogram if possible before your vaccine. However, if you are already overdue for your screening exam or cannot reschedule within the next few months, keep your screening mammogram appointment and keep your COVID-19 vaccination appointments. Regular screening mammograms ensure that breast cancer can be detected as early as possible. Both are very important to ensure that you stay healthy. It's especially important to keep your mammogram appointment if you are significantly overdue for screening. Remember: annual screening saves the most lives.

■ What should I tell the technologist on the day of my screening mammogram?

Notify your mammography technologist if you have received a COVID-19 vaccine. Tell her when you received the vaccine, and which arm the vaccine was given. State whether it's your first or second dose. This information will help the breast radiologist interpreting your screening mammogram.

■ What if I have other breast problems?

If you have any changes in your breast or underarm, such as pain or a lump, contact your medical provider. The guidelines above are only for women with no breast symptoms who are scheduled for a COVID-19 vaccine.

